

Ca



Product information

Lebosol®-Calcium

EC FERTILISER | calcium chloride solution | p 1.32 | pH 7.5

Nutrients (w/w)	g/l
16.7% water-soluble calcium oxide	220

Crop	Objective	Recommendation
In crops insensitive to chloride	To provide calcium	5 - 10 l/ha with foliar fertiliser in at least 500 l water. Upon application with backpack sprayer 1%.
Potatoes	Tuber and skin quality, improvement in storage life	3 times 5 l/ha from row closure
Strawberries	Fruit firmness, storage stability, resilience	3 - 4 times 5 l/ha between fruit set and harvesting
Pome fruit	Fruit firmness, storage stability, resilience	4 - 6 times 5 - 10 l/ha from walnut size to harvesting
Stone fruit	Fruit firmness, storage stability, resilience	2 - 5 times 5 - 10 l/ha between fruit set and harvesting
Soft fruit	Fruit firmness, storage stability, resilience	3 - 4 times 5 l/ha between fruit set and harvesting
Dessert grapes	Fruit firmness, storage stability, resilience	3 times 5 l/ha from pea size to harvesting
Wine grapes	Fruit firmness, storage stability, resilience	3 - 5 times 5 l/ha from pea size to harvesting
Fruiting vegetables	To prevent blossom end rot, fruit firmness, storage stability, resilience	3 - 4 times 5 l/ha from fruit set
Cabbages, leafy vegetables and bulb vegetables	To prevent tipburn and leaf tip necrosis, increase in storage life, resilience	3 - 4 times 5 - 10 l/ha once sufficient leaf mass has developed
Asparagus	Increase in storage life, resilience	2 - 4 times 5 - 10 l/ha once sufficient leaf mass has developed
Root vegetables and tubers	Fruit firmness, storage stability, resilience	3 - 4 times 5 - 10 l/ha once sufficient leaf mass has developed
Medicinal plants, scented plants and spice plants	Fruit firmness, storage stability, resilience	2 - 4 times 5 - 10 l/ha once sufficient leaf mass has developed or from the end of flowering to harvesting

You can find more information on the hotline: +49 (0) 63 28-9 84 94-80 or on our website www.lebosol.de.